


Antifungals and Antivirals



Across

2. Infectious EBV is also known as this
3. Posaconazole is contraindicated with _____
4. Brand name for Posaconazole
5. side effect of Voriconazole
6. the only neuraminidase inhibitor that does not require renal adjustment
8. this is considered fluconazole resistant (include the .)
10. associated with causing Thrombocytopenic Purpura
11. 5FC shouldnt be used as _____
13. warning associated with adamantanes
14. Cryptococcus Neoformans is a _____
18. Acyclovir resistant HSV lesions are treated with _____ until the lesions heal
24. This antiviral is a vesicant, and should ideally be given via central line
26. CMV is a type of _____ virus
27. should be used with 5FC for treatment of invasive cryptococcal meningitis or Candida infections
28. The most commonly identified cause of Viral encephalitis
29. all Azole antifungals have this adverse effect except isavuconazonium (no spaces)
31. Generic name of Mycelex
32. HSV2 is associated with _____ disease

35. Amphotericin B (conventional) can cause what if overdosed?
37. Echinocandins can cause _____ mediated symptoms
38. The only azole antifungal that requires renal adjustment
39. First line treatment for Candida glabrata or Candida Krusei
40. frequency of which echinocandins are given
41. can be used for pain caused by shingles
42. This antifungal needs to be taken with a fatty meal to increase absorption and avoid GI upset
44. Boxed warning for Ganciclovir

Down

1. Zygomycetes are also known as _____
7. Tamiflu can be given to anybody _____(age)_____ and older (no spaces)
9. Azole antifungals decrease synthesis of this _____
12. Duration of treatment (days) for a patient diagnosed with Herpes zoster
15. This antifungal has a 1:1 IV:PO ratio
16. Amantadine is not recommended due to _____
17. ketoconazole and itraconazole cannot be taken with this _____
19. brand name of Flucytosine
20. HSV1 is associated with _____ disease

21. form of Acyclovir used for herpes simplex labialis
22. this drug (brand) requires a filter for administration
23. acyclovir is dosed based on _____ body weight
25. The higher end of the therapeutic range for Voriconazole
30. This antifungal should be given PO if the pts CrCl is 50ml/min
33. Brand name of Caspofungin
34. Brand of Liposomal Amphotericin B
36. a Griseofulvin Contraindication
43. The lower end of the therapeutic range for Voriconazole