


Name: _____

Date: _____

Cardiac Unit Competency



Across

2. A collection of blood outside of a blood vessel.

6. A health-risk behavior that is one cause of heart disease.

8. A reading of <65 of this clinical measure is critical and requires RRT activation.

15. A common complication of TAVR surgery.

18. Cardiac patients have higher risk of _____ than most of the general population.

19. This type of pacing improves symptoms of heart failure by coordinating both ventricles.

20. Beta Blockers may mask this condition.

21. A symptom of cardiac tamponade

22. Patients with which type of MI can wait to go to the cath lab if anticoagulation cardiac demand reduction is given?

24. Cardiac ablation comes with a risk for this fatal complication.

25. Vital signs are imperative before and after giving each dose of this medication.

26. What drug category is required for 4-6 weeks after placing a bare metal stent?

Down

1. Sotalol has a black box warning for tachycardia due to?

3. RT must be notified for increasing _____ requirements.

4. Over time, this condition leads to heart failure

5. A type of communication that aides in preventing sentinel events.

7. Antithrombotic medication may be necessary for a few weeks prior to this procedure.

9. Having a patient lean forward may be helpful in differentiating chest pain of an MI from _____.

10. A condition that elevates risk for adverse effects of epidural analgesia.

11. A type of ablation often performed along with open heart surgery.

12. A type of ablation performed for dysrhythmias

13. A device used to decrease risk of stroke in patients with afib.

14. It is important to check and normalize potassium and magnesium prior to giving _____.

16. A medicine commonly used to treat pericarditis.

17. Equipment used to keep the airway open, improving gas exchange and reducing or eliminating the need for intubation.

23. Prior to sending a patient for an ablation, which medication should be given?