


Chapeter 40, 41, and 42



Across

3. Children with cystic fibrosis take this with meals and snacks
 4. This type of rice is not high in fiber
 6. A patient should not receive iodine contrast if they are allergic to this?
 11. Normally pressure is higher on this side of the heart
 14. This medication should be used for an asthma attack
 15. This defect causes decreased pulmonary blood flow
 16. This is the number of hours a child with streptococcal pharyngitis must stay home from school after starting abx therapy
 17. What is a s/s to administer epinephrine after accidentally eating a food you are allergic to

19. If the child vomits with this many minutes after administering digoxin, it is alright to give a 2nd dose.
 22. What do you do if the dressing is soaked with blood after cardiac cath?
 23. With a septal defect blood will flow from left to right
 24. This is a high fiber food?

Down

1. What is the leading cause of death after a heart transplant?
 2. A child's extremity should be in this position after cardiac catheterization
 5. This should be avoided with a child who has a diagnosis of Hirschsprung?
 7. This is a sign of choking in a toddler
 8. What do you add to formula to minimize reflux?
 9. This is a sign of digoxin toxicity

10. The pulse is determined by this rate when administering digoxin
 12. This is a heart defect present at birth
 13. What snacks are appropriate for children with cystic fibrosis?
 18. This is the best prevention of spreading gastroenteritis
 20. A cardiac cath cannot be performed if a child has this.
 21. Blood flows from an area of higher pressure to an area of