


Child & Family Rotation



Across

3. Deficiency of glucose in the bloodstream

8. Loss of more than 500-1000mL of blood within the first 24 hours following childbirth

10. A common condition in newborns when there is too much bilirubin in the blood

12. Average _____ for newborns is 36.5 - 37.2

13. Health professionals who provide primary care to women and their babies during pregnancy, labour, birth, and the postpartum period

14. After 24 hours a newborn requires a/an _____ before discharge

16. Intervention to promote maternal-newborn bonding, as well as elevation of newborn temperature after baths

17. Recommended method of newborn feeding

18. The _____ is midline and palpable halfway b/w the symphysis pubis and the umbilicus immediately after delivery

Down

1. A c-section patient would require a _____, and a vaginal delivery patient may not

2. respect for patient's unique background or way of life demonstrates _____

4. Solution made from human blood plasma that is given to Rh negative women

5. a large part of working on the postpartum unit involves _____. Some examples of this are breastfeeding frequency, latching, ambulation, self-care, etc.

6. Most common treatment for reducing high bilirubin levels that cause jaundice in a newborn

7. A woman without diabetes develops high blood sugar levels during pregnancy

9. Delivery of a baby through a surgical incision in the mother's abdomen and uterus

11. A rubella nonimmune mother would require a _____ injection

15. common bacteria which are often found in the vagina, rectum, or urinary bladder of women and can lead to newborn complications if not treated