
Name:	Date:
name:	Date:

## Drug Doses



## Across

- 2. Initial dose 0.4 2 mg, but has been given up to 10 mg in the field
- **4.** Your pt has hives on their chest and they itch. What do you give?
- 8. Dose is 1 meq/kg
- 13. We give it as a drip for Bradycardia and/or Hypotension 14. Initial dose for narrow complex tachycardia 0.25 mg/kg 6. A contraindication of giving

17. Pain management initial dose is 2-5 mg

**20.** Solution is 25% when given to children less than 12 years old

## Down

- 1. Given for seizures if pt is allergic to Versed
- **3.** The dose is 1 mg 1:10,000 solution and given for cardiac arrest
- 5. Adult and Pediatric dose in 2.5 mg/3 ml
- this medication is a BP of 90 sytolic
- 7. Given for Hyperkalemia

- **9.** Last drug given for refractory asthma
- 10. Mixed with Albuterol during first administration
- 11. Another name for Lasix
- 12. Given PO for Chest Pain patients
- 15. Pediatric dose is 5 mg/kg given over 20-60 minutes
- **16.** Pain medication given with a dose of 1 mcg/kg
- 18. Can't be given to children less than 6 years old
- 19. Pediatric dose for Asthma is 2 mg/kg