


GI Meds Continued...



Across

4. Sucralfate can interfere with absorption of phenytoin, digoxin, warfarin, and ciprofloxacin. The nurse should give these meds at least _____ apart to minimize interaction.

9. When evaluating effectiveness of Probiotics, nurses hope to assess a _____ frequency of manifestations of hiatal hernia such as heartburn, belching, and dysphagia.

11. MOA: muscarinic and histaminergic receptors in nerve pathways that connect the inner ear and VC are blocked.

14. Sucralfate does not have _____ effects because it is not absorbed.

17. Absorption of ketoconazole, itraconazole, and atazanavir is _____ when taken concurrently with proton pump inhibitors.

18. When measuring the effectiveness of Prostaglandin E analogs, nurses should assess for the frequency of GERD, absence of GI bleeding, and presence or healing of _____.

19. Prokinetic agents are effective when the nurse assess that there is an _____ of nausea and vomiting.

20. Nurses should remind patients with Peptic ulcer disease to avoid drinking _____ and to notify the provider if there is any indication of GI bleeding

21. The therapeutic use of Cannabinoids is to control CINV and to _____ appetite in patients that have AIDS

22. _____ laxatives result in stimulation of intestinal peristalsis

24. What is the prototype for Antacids?

25. Antacids containing sodium can cause fluid _____. Teach patients with hypertension or heart failure to avoid antacids that contain sodium.

Down

1. Dimenhydrinate (a antihistamine) can be administered PO, IM, or _____

2. When administering Dopamine antagonist, nurses need to consider clients on antihypertensives and monitor closely for _____.

3. Magnesium Compounds can cause _____.

5. Iron and _____ can alter the absorption of sulfasalazine.

6. Use cannabinoids cautiously in those patients with _____ disease

7. Aluminum and calcium compounds can cause _____.

8. When administering Lubiprostone, we know it is effective when there is a relief of constipation or diarrhea, or if the therapeutic use is _____.

10. Pantaprazole is administered to patients _____.

12. Irritation at the injection site after administering Pantoprazole, there is a risk for _____. The nurse should monitor site closely for inflammation.

13. Advise clients that CNS depression is more likely and to avoid activities that require mental _____.

15. Dronabinol (a cannabinoid) is administered by _____.

16. Antacids are used to treated PUD and _____ by promoting healing and relieving pain.

23. To prevent constipation when taking Sucralfate, the nurse should encourage clients to increase intake of fiber and _____ at least 1500 mL of fluids if fluids are not restricted.