
HELLP SYNDROME



<u>Across</u>

1. _____ is the breakdown of red blood cells.

8. __ pain is found in 65% to 90% of patients with HELLP Syndrome. **11.** Placental _____ is a potential complication of HELLP Syndrome. 14. The clinical course of women with true HELLP Syndrome is often characterized by a _ deterioration in the maternal condition. 16. One of the three chief abnormalities found in the presence of HELLP Syndrome is elevated enzymes. 17. _ _ sulfate is administered to prevent seizures. 18. HELLP Syndrome sometimes

presents during the _____ period.

19. _____ edema is a complication sometimes associated with HELLP Syndrome.

20. Laboratory abnormalities in HELLP Syndrome typically worsen after _____ and then begin to resolve 3 to 4 days after.

Down

2. _____ is a serious blood clotting complication of HELLP.

 Prematurity, _____, and thrombocytopenia are neonatal complications of HELLP Syndrome.
_____ women are at higher risk for HELLP Syndrome.

5. Which race is at higher risk for HELLP Syndrome?

6. Serious hepatic manifestations of HELLP include infarction, hemorrhage, and _____.

7. Most often, HELLP Syndrome develops in the _____ trimester of pregnancy.

9. _____ is a rare complication of HELLP Syndrome.

10. The cause of HELLP Syndrome is

12. Between 38 and 93 percent of HELLP patients receive some form of _____ product.

13. It is theorized that HELLP Syndrome is a variant of _____.

15. ______ failure is a possible complication of HELLP Syndrome