


IBS/IBD/GERD/PUD



Across

1. excessive fat in the stools due to malabsorption
3. generic of Uceris, used to help treat ulcerative colitis
4. Blood in stool
5. ADR of anti-histamine's, dizziness, spinning room
6. Generic name of Tagamet that has many drug interactions
8. IBS-C in women, OI-Constipation
11. Brand only, PPI
18. Side effect of GERD, ADR of Imodium
22. _____ tenderness, indicates severe to fulminant UC
23. Go to antacid, active ingredient in TUMS
25. When there are drug interactions with antacid use, separate antacid by ____ hours
28. Gives greatest symptom relief for GERD + highest healing rates

31. Malignant gastrinomas occur in 30-50% of patients with this syndrome
32. Not recommended for acid reflux, dosed QID, forms viscous adhesive on ulcer crater
33. Omeprazole and sodium bicarb
34. 250/500 mg capsules that are effective in entire GI tract
35. Weakening of bone, extra intestinal manifestation
36. used for IBS-D and hits Mu receptors
37. Food that decreases LES pressure

Down

2. A dangerous complication of GI ulcers
7. antidiarrheal that is over the counter
9. use of loperamide may cause this condition
10. classic symptom of PUD
12. Protective factor for PUD
13. Medication that is an irritant to mucosa in GERD

14. Initial treatment for ulcerative colitis
15. Prokinetic agent, increases LES pressure
16. Generic name for Sandostatin (inhibits gastric acid and releases gastrin)
17. An independent risk factor for stress-related mucosal bleeding
19. Barrier to effective treatment of H. pylori infection
20. emergency use in IBS-C and CIC
21. Mucus secreting cells in intestinal lining
24. lifestyle modification for nocturnal symptoms of GERD
26. PPI's are dosed 30-60 mins before which meal?
27. Drug used for prevention of NSAID ulcers, but contraindicated in pregnancy
29. ____ weeks of PPI is the choice for symptomatic relief and healing of erosive esophagitis
30. Food may relieve the pain when dealing with a ____ ulcer