


Name: \_\_\_\_\_

# Janssen Drugs / Indication



## Across

1. Indicated for several disease states such as Crohn's Disease, RA, Ulcerative Colitis, etc.
2. Active psoriatic arthritis (PsA) Active ankylosing spondylitis (A)
5. Indicated for the treatment of adult patients with active psoriatic arthritis, and Crohn's Disease.
9. Indicated for the treatment of patients with unresectable or metastatic liposarcoma or leiomyosarcoma who received a prior anthracycline-containing regimen.
12. In combination with other antiretroviral agents (ARVs), is indicated for the treatment of human immunodeficiency virus (HIV-1) infection in patients.
13. Indicated for the treatment of human immunodeficiency virus type 1 (HIV-1) infection in ARV treatment-experienced patients ages 6 years and older,

15. Indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus.
16. Indicated for the treatment of adults with moderate to severe plaque psoriasis
17. Indicated in combination with other antiretroviral agents for the treatment of human immunodeficiency virus (HIV-1) infection
18. Indicated to reduce the risk of stroke and systemic embolism in patients with nonvalvular atrial fibrillation (AF).
19. Indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus when treatment with both canagliflozin and metformin is appropriate.

## Down

3. Indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus when treatment with both canagliflozin and metformin is appropriate.

4. Indicated for the treatment of human immunodeficiency virus type 1 (HIV-1) infection in antiretroviral treatment-naïve patients 12 years of age and older
6. Indicated for the treatment of patients with Multicentric Castleman's Disease
7. Moderately to severely active rheumatoid arthritis (RA) in combination with methotrexate
8. Indicated for with metastatic castration-resistant prostate cancer (mCRPC) with metastatic high-risk castration-sensitive prostate cancer (CSPC)
10. Indicated for the treatment of patients with non-metastatic castration-resistant prostate cancer.
11. Indicated for the treatment of anemia due to chronic kidney disease (CKD),
14. For the treatment of patients with newly diagnosed multiple myeloma