

Name: _____ Date: _____

Kidney Transplant Medications

- | | |
|---|-------------------------------|
| 1. An antifungal medication that is taken four times a day | A. Headaches or tremors |
| 2. An antiviral medication that is taken to protect you from certain viruses like CMV | B. Prograf / Tacrolimus |
| 3. A medication that suppresses your immune system and protects your kidney from rejection. We check levels of this medicine. | C. Nystatin |
| 4. An antibiotic that is used to prevent/treat urinary tract infections and PCP pneumonia | D. Nausea, vomiting, diarrhea |
| 5. An immunosuppressant medication commonly taken in addition to Prograf. | E. Bactrim |
| 6. Cellcept or MMF may cause these side effects | F. Cellcept/ MMF |
| 7. It is important to check your Prograf levels, because high levels can cause these symptoms _____ | G. Low magnesium |
| 8. Prograf can causes this problem with your electrolytes | H. Valcyte / valganciclovir |
| 9. Valcyte can cause your white blood cell count (infection fighting cells) to be _____ | I. Rejection |
| 10. It is important to take your Prograf every 12 hours and get your levels checked, because low levels can place your kidney at risk for _____ | J. Low |
| 11. This is a medication that is commonly used to treat high blood pressure and may cause swelling | K. Amlodipine |
| 12. Prograf (a transplant medication) and Lisinopril (a high blood pressure medication) can sometimes cause ____ potassium levels. | L. High |