
name.	Name:		Date:	
-------	-------	--	-------	--

Labor and Delivery

Across

- 2. The fetus is in a longitudinal lie with the buttocks or feet closest to the cervix
- **6.** Secreation of milk from the breast
- **9.** The fluid surrounding a fetus
- **11.** The normal discharge from the uterus after childbirth
- **12.** The lower part of the uterus that dilates as labor progresses
- **16.** Used to determine the position of a fetus inside the woman's uterus
- **18.** A synthetic form of oxytocin, used to induce labor
- **19.** Shortening of the uterine muscles occurring at intervals before and during childbirth





Down

- 1. Cervical thinning
- **3.** A surgical cut made at the opening of the vagina during childbirth
- **4.** The procedure by which the amniotic sac is deliberately ruptured
- **5.** A decrease in the fetal heart rate below the fetal baseline
- 7. The sensation that a pregnant woman feels when the baby drops
- **8.** The fluid-filled sac that contains and protects a fetus in the womb; membranes

- **10.** The act of giving birth; childbirth
- **13.** The opening of cervix as it is pulled upward and fetus is pushed downward
- **14.** The top portion of the uterus
- **15.** A situation at childbirth where the fetus is in a longitudinal lie and the head enters the pelvis first
- **17.** The location of the presenting part of the fetus in the birth canal