


Name: _____

Date: _____

Medication Administration: Medication List



Across

- 2. Given SQ 2 inches away from umbilicus
- 3. Loop diuretic
- 5. Monitor electrolytes, daily weight, BP, AP, I/O, lung sounds, edema
- 6. Antidote: Protamine Sulfate
- 7. Side Effect: dark, tarry stools
- 8. Bronchodilator
- 14. antianginal
- 21. Side Effects: insomnia, wt loss, heat intolerance, tachycardia
- 22. Antiarrhythmic
- 24. Monitor for bleeding: look at platelets
- 25. antidote: Vitamin K
- 26. Non-opioid analgesic, antipyretic
- 27. anticoagulant
- 31. Iron supplement

- 33. Side Effect: Severe Headache
- 34. Cloudy, roll
- 35. Electrolyte
- 36. Tx tremors and rigidity sx
- 37. Symptoms: Polyuria, Polyphagia, Polydipsia
- 38. Antidote: Protamine Sulfate

Down

- 1. Side effect of Morphine and Hydrocodone/APAP
- 4. Toxic doses may cause renal toxicity
- 9. Monitor K+ levels, may alter rhythm of the heart
- 10. Side effect of Morphine & Hydrocodone/APAP
- 11. Stool Softener
- 12. Given SQ, is clear and can be given IV
- 13. Non-opioid analgesic, antipyretic

- 15. Treats hypothyroidism
- 16. Medication to be available for Morphine & hydrocodone/APAP
- 17. toxicity with hypokalemia--anorexia, fatigue, confusion, bradycardia
- 18. antiplatelet -prevents platelet aggregation (platelets coming together to form a clot)
- 19. Check AP before admin.--HOLD below 60 BPM/ or above 100 bpm
- 20. Toxic doses may cause hepatotoxicity
- 23. anticoagulant
- 28. Anticoagulant given SQ in anterolateral and posterolateral abd
- 29. Watch for GI Bleeding
- 30. Monitor for bleeding using PTT
- 32. Side Effect: Tinnitus