


# Moderate Sedation and Capnography



## Across

3. A patient with poorly controlled diabetes can be expected to have an ASA class of \_\_\_\_\_
4. Anesthesia will be consulted for airway management and monitoring for a patient with an ASA \_\_\_\_\_ class.
6. Any patient who has IV \_\_\_\_\_ at any point during the administration of sedation will require an additional 2 hours of observation and monitoring.
7. During the procedure, Vitals, RASS, \_\_\_\_\_, and cardiac rhythm will be documented every 5 minutes

11. A \_\_\_\_\_ by team members regarding the appropriateness of the ASA classification assignment must be reached prior to initiating the procedure.
12. All patients will have a pre and \_\_\_\_\_ Aldrete Score assigned.
13. Moderate sedation administration is limited to the following areas: endoscopy, CT, Interventional Radiology, critical care units, ED, Surgical Services Pre-op, Open Heart, Care Unit, and \_\_\_\_\_.
14. Reflex \_\_\_\_\_ from a painful stimuli is not considered a purposeful response.

## Down

1. The team is responsible for performing a time out \_\_\_\_\_ the administration of medications
2. \_\_\_\_\_ must be held for 4 hours prior to moderate sedation
5. What is the reversal agent for Versed?
8. The reversal agent for Fentanyl is \_\_\_\_\_,
9. Moderate sedation is a drug induced depression of consciousness during which patients cannot be easily aroused, but \_\_\_\_\_ purposefully.
10. Anesthesia does not need to be consulted for moderate sedation if the patient is on this device.