
Name:	Date:

NEWBORN



<u>Across</u>

- 2. blueness of hands feet of newborn after delivery due to high hemoglobin level-normal phenomenon
- **5.** occurs when you stroke the sole of the foot and big toe dorsiflexes and toes flare; Birth ends before walking
- 7. unossified spaces or soft spots on the cranium of an infant; protect head during delivery and allow for brain growth in first 1 1/2 yrs
- **8.** infant draw up legs and arms fan out & then come toward midline when this is done (startled like if hands clapped near infant or crib jarred suddenly)

- **9.** prancing movements of legs seen when infant is help upright on the examining table birth to 4-5 months
- **11.** surgical removal of foreskin on penis
- 13. cheese like substance that covers skin of newborn; made of cells and glandular secretions; thought to protect the skin from irritation and the effets of a watery environment in utero

Down

1. milia type lesions on the midline of the hard palate caused by collection of epithelial cells

- **3.** dark blue or slate grey discolorations found mostly in lumbosacral area; caused by melatonin deposits in dark skinned persons; gradually disappears in few years
- **4.** causes infant's head to turn in direction of anything touching cheeks anticipating food; use to help with breast feeding; Birth to 4 months
- **6.** dark, thick, sticky, first stool of baby
- **10.** fine hair covering baby at birth; disappears during first week of life; more evident in premie babies
- **12.** white pinpoint pimples caused by obstruction of sebaceous glands seen on nose & chin of infant