


# NSG 106 Test 3 & 4 Meds



**Across**

- 3. Generic name for Flexeril
- 8. 'LOL anti-hypertensive drug #2
- 9. What 2 drugs are used to treat parkinson's disease?
- 11. ACTION: Block the sympathetic nervous system. If it's blocked everything decreases so get low BP and decreased pulse. USE: Treat hypertension, angina, cardiac arrhythmias.
- 12. ACTION: Increases the strength of heart contraction USE: Treat congestive heart failure SIDE EFFECTS: Bradycardia, dizziness, nausea or vomiting, halos before eyes
- 13. Muscle Relaxer that can also be used for anti-anxiety

- 14. Non steroidal anti-inflammatory drugs
- 15. Common Side Effects: Sedation, disorientation, coma, constipation, respiratory depression
- 17. 'LOL anti-hypertensive drug #3
- 18. Can cause Reyes Disease in children

**Down**

- 1. ACTION: Mimics the sympathetic nervous system and increases everything (especially vital signs). Also stimulates beta 2 receptors in the lungs which produce dilation of the bronchial tubes, increasing air flow to the lungs.
- 2. ACTION: Block histamine receptors reducing the effects of histamine release. What is the sedating drug name?

- 4. ACTION: Cause K and Na to be excreted from the Loop of henle in the kidneys. Wherever K and Na go water will follow because water follows salts and K and Na are salts. This will cause K, Na and water excretion (urine)
- 5. Acetaminophen brand name
- 6. NURSING IMPLICATIONS: Usually administered via inhalation.
- 7. 'LOL anti-hypertensive drug
- 10. It is important to assess this function when patient is taking Acetaminophen
- 16. Causes ringing in the ears after access dosing