
Name:	Date:	
10.11101	_ ~	

Parkinson's Disease



Across

- **3.** A peripheral decarboxylase inhibitor used in combination with levodopa to reduce adverse effects of levodopa
- **4.** One of the four classic symptoms of Parkinson's disease
- ${\bf 5.}$ Newest antiparkinson drug approved by FDA in 2008
- **6.** Mean age of onset of Parkinson's disease (2 words)
- 16. Slowness of movement
- 17. The biologic precursor of dopamine that can penetrate into the $\ensuremath{\mathsf{CNS}}$
- **19.** One example of newer nonergot nondopamine dopamine receptor agonists
- **20.** One drug under catechol ortho-methyltransferase (COMT) inhibitors category
- ${\bf 21.}\ \mbox{An example}$ of an indirect-acting dopamine receptor agonist
- **22.** Type of tremor seen mostly at rest and less severe during voluntary activity (2 words)

- **24.** How many millions of Americans are affected by Parkinson's disease?
- **25.** This phenomenon is a gradual worsening of Parkinsonian symptoms as a patient's medications begin to lose their effectiveness (Hyphenated)
- ${\bf 27.}~{\bf Absence}$ of psychomotor activity resulting in mask-like facial expression
- **28.** Destruction of this part of the brain by Parkinson's disease leads to dopamine depletion (2 words)
- **29.** Abnormal muscle tone leading to impaired or abnormal movements
- **30.** Levodopa and carbidopa are both contraindicated in this type of glaucoma (2 words hyphenated)

Down

- 1. An excitatory neurotransmitter
- 2. An inhibitory neurotransmitter
- 7. A major effect of nonselective MAOIs when they interact with tyramine-containing foods (2 words)
- 8. This COMT drug is contraindicated in liver failure
- **9.** An irregular, spasmodic, involuntary movements of the limbs or facial muscles

- 10. Enzyme that causes breakdown of catecholamines in the body (2 words)
- 11. Second most common neurodegenerative disease
- **12.** An antiviral drug considered a dopamine modulator that is also indicated for treatment of moderate Parkinson's disease
- **13.** An anticholinergic drug used for Parkinso's disease
- 14. Type of rigidity seen in Parkinson's disease
- **15.** A drug under ergot direct-acting dopamine receptor agonist
- 18. Shakiness of the extremities seen mostly at rest
- **23.** Drugs that exert their antiparkinson effects before the nerve synapse
- **26.** Symptoms of Parkinson's disease do not appear until how many percent of dopamine in the substantia nigra has been depleted?