


Name: _____

Date: _____

Pediatric emergencies



Across

2. All cardiac arrest will be worked for at least ____ minutes on scene prior to the transportation of the patient
 6. Administer _____ (Benadryl) 1mg/kg slow IVP for mild allergic reaction
 11. When the skin is visibly more pale in certain areas of a child
 12. Aside from extrication, on scene time should be no greater than ____ minutes
 14. Do not use _____ splint if a pelvic fracture is suspected
 15. approach a child _____ to _____. opposite of an adult
 17. enables ems to treat and give rapid dosing for the critically ill pediatric

19. When the skin appears blotchy, web like pattern

20. Sound of lower Airway-acute bronchospasm

Down

1. Administer _____ (Midazolam) 0.1mg/kg for a Seizure
 3. Upper Airway noise
 4. 1-4 years, a burn on a child head is _____
 5. Glucose levels are <70 with sign/symptoms
 7. _____ Control Center 1800-222-1222

8. Administer _____ 1:1000- .01 mg/kg IM. max dose of .3 in severe allergic reactions

9. Administer _____ if unable to establish iv access with a glucose below 70

10. In children > 1 year of age, Administer _____ 1-2 mcg/kg SLOW IVP, repeated once.

13. a seal-like cough/bark

16. Suspected Opiate or Opioid Overdose. Administer _____ 0.1 mg/kg max dose of 0.5

18. Defibrillate at joules/kg for a shockable rhythm

Word Bank

ten	Versed	Traction	Narcan	Pallor
Toe to head	Diphenhydramine	Epinephrine	nineteen	two
Hypoglycemic	Mottling	Handtevy	twenty	Croup
Fentanyl	Stridor	Wheezing	Poison	Glucagon