


Name: _____

Return completed puzzle to PICC Team by Nov. 30th 2018.



Across

2. the Central Venous Catheter Discontinuation _____ will be used for removal of temporary Central Lines and will be returned to the nurse manager once completed.
7. Patients _____ with a Midline in place must also have a referral for home health/infusion clinic/to a health care facility.
8. IV sites and infusions should be assessed a _____ of every 2 hours.
9. Swab Caps must be in place at least 5 minutes in order to _____ the need to scrub for 15 seconds with an alcohol prep pad.
11. Site _____ includes signs and symptoms of phlebitis, infiltration, leaking of IV fluid, sensation changes, and intactness of dressing.

12. Acceptable PICC tip locations are the _____ or Cavo-Atrial Junction.

13. _____ that are not appropriate for peripheral IV's are also not appropriate for Midline catheters, as a midline is not a central line.

Down

1. Prior to Venipuncture, sites should be scrubbed for a minimum of 30 seconds with 2% _____.
3. The _____ of the PICC line should be measured and compared to the external length at the time of placement every shift for inpatients and every visit for outpatient infusion areas.
4. The _____ method should be used to clear incompatible medications/IV's from secondary intermittent IV tubing.

5. Multiple _____ attempts cause patient pain, delay treatment, limit future vascular access, increase cost, and increase the risk of complications.

6. If the patient is admitted with a previously placed PICC line, the PICC location must be _____ with chest x-ray upon admission to the hospital, prior to administering medications or IV fluids.

10. _____ disconnecting and reconnecting IV tubing for activities such as showers, baths, etc.