


Name: \_\_\_\_\_

# SEPSIS HEROES



## Across

3. THE \_\_\_\_\_ RELAY IMPORTANT INFORMATION TO IMPROVE HEALTHCARE IN A FACILITY.
5. For sepsis hypotension, the recommended crystalloid fluid bolus is 30 mls per \_\_\_\_\_.
6. Clinical documentation \_\_\_\_\_ query providers for additional documentation so that the record is clearer and the reimbursements are appropriate.
13. Blood \_\_\_\_\_ must be drawn prior to antibiotics to give the best results.
19. IF LACTIC ACID RESULT IS >2, WHAT MUST BE DONE WITHIN 2-3 HOURS?
20. If the boluses aren't bringing the systolic blood pressure back up to >90 or MAP >65, best vasopressor is?
21. WHO MIGHT CALL A SEPSIS ALERT?
22. Family members can save lives by recognizing \_\_\_\_\_ of sepsis.
24. Emergency Medical Service (acronym) are Heroes Every Day
25. It is important for the community to understand that if they THINK sepsis, they should \_\_\_\_\_ sepsis when bringing a family member to the ED.
26. FLUID \_\_\_\_\_ are important to reperfuse organs when side-effects of sepsis has leaked fluids out of the vessels.
28. Chemistry lab personnel contact nurse if initial lactic acid is \_\_\_\_\_.

29. Clotting problems may occur during and after septic conditions therefore it is best that patients follow-up as directed with their \_\_\_\_\_.

30. SEPSIS HEROES RARELY GET A RESTROOM \_\_\_\_\_.

32. fast heart rate, fast respirations, high or low temperature, high or low white blood count, or Bands >10% are signs of systemic inflammatory response syndrome or \_\_\_\_\_.

33. After a patient goes into Septic Shock and has fluid boluses given, it's important for physician, nurse practitioner or physician assistant to perform a \_\_\_\_\_ exam and for Nurses to obtain and document frequent Vital Signs.

34. Common side effect of sepsis within the next year is a \_\_\_\_\_.

35. Timely \_\_\_\_\_ lowers risk of mortality

36. When patient has 2+ SIRS, infection and organ dysfunction, they have \_\_\_\_\_ sepsis.

## Down

1. Carries a cell phone, a prescription pad, and sometimes the weight of the world.

2. If we provide evidence-based treatment swiftly for sepsis patients, we are reducing risk of \_\_\_\_\_.

4. First greets and assesses patient and directs care in ED.

7. Supervisors, managers, directors and Executives are Sepsis Heroes even when not at the \_\_\_\_\_.

8. Vehicle with sirens and lights to get patients swiftly to hospital when have symptoms of sepsis.

9. These Heroes who wield a scalpel, can sometimes cut away the source of infection.

10. Administrative Personnel support our Heroes in many roles in \_\_\_\_\_.

11. CODE \_\_\_\_\_ IS CALLED BY ANYONE WHO FINDS A PATIENT WITH 2 SIRS, A SUSPECTED INFECTION AND ORGAN DYSFUNCTION

12. Nurses who follow Sepsis Protocol are their patient's \_\_\_\_\_.

14. \_\_\_\_\_ can show if organs are perfusing.

15. SHOULD BE ADMINISTERED WITH A PRESSURE BAG TO REPERFUSE FOR HYPOTENSION OR LACTIC ACID >4 MMOL/L

16. INPATIENT PHYSICIANS CAN REDUCE SEPSIS READMISSIONS IF THEY HAVE PATIENT FOLLOW-UP WITHIN 7 DAYS AND HAVE HOME HEALTH \_\_\_\_\_ DAY AFTER DISCHARGE.

17. These technicians swiftly draw their samples and send them to the lab.

18. LEVELS >2 ARE INDICATIVE OF ORGAN DYSFUNCTION FOR CREATININE, LACTATE AND \_\_\_\_\_.

23. WHO COMMUNICATES A SPOT ALERT TO NURSE?

27. TAKE CARE OF SEPSIS PATIENTS FROM THEIR HOME TO THE HOSPITAL

31. Broad \_\_\_\_\_ Antibiotics are best initially, until blood cultures come back and the first dose should be administered within one hour of presentation.

## Word Bank

PHLEBOTOMISTS  
MONITORTECH  
AMBULANCE  
norepinephrine  
healthcare  
kilogram

BILIRUBIN  
ANTIBIOTICS  
Focused  
REPEAT  
PARAMEDICS  
EMS

MORTALITY  
SIRS  
severe  
TRIAGE NURSE  
say  
spectrum

surgeons  
SEPSIS  
Doctor  
Heroes  
Lactate  
Specialists

BREAK  
cultures  
symptoms  
VISIT  
Physician  
FLUIDS

BOLUSES  
bedside  
stroke  
EDUCATORS  
elevated  
NURSE