


STIs in Pregnancy



Across

2. Which treatment for chlamydia is contraindicated during pregnancy and could reduce the effectiveness of oral contraceptives?
4. What is the most common treatment for GBS in labor?
6. Which infection typically has no manifestations in females yet can lead to tubal scarring and lead to PID?
9. What is the best way to prevent cytomegalovirus?
10. Which infection is caused by consumption of raw or undercooked meat or handling cat feces?
11. Which antibiotic is the medication of choice for the neonatal prophylactic treatment against gonorrhea and chlamydia?

13. Which common STI manifests with small cauliflower-like warts in the genital area?

14. During pregnancy, which infection is treated to prevent oral thrush in the neonate?
15. Which bacterial infection is tested between 35 to 37 weeks gestation using a vaginal and rectal swab?
17. A combination of sulfadiazine and pyrimethamine, although potentially harmful to the fetus, is used for the treatment of which infection?
18. Which medication should the nurse expect a provider to order for the treatment of gonorrhea?

Down

1. Black, Hispanic, and other racial/ethnic minority groups in the United States are disproportionately affected by this STI.

3. Which is the recommended way to deliver for clients in labor who have active genital herpes lesions or early findings of an impending outbreak?

5. How many stages are there to Syphilis?
7. Which STI is manifested in females with a yellow-green, frothy vaginal discharge with a foul odor?
8. Which STI is the most commonly reported in American women?
12. Which TORCH infection manifests with painful blisters and tender lymph nodes?
16. Which vaccine to protect against a TORCH infection is not given during pregnancy?