


Name: _____

Date: _____

Sepsis and PICC



Across

1. Sepsis is SIRS with documentation of a presumed or ___ infection
4. In the ___ bundle, repeat lactate if initial lactate > 2
6. Septic shock is sever sepsis with lactate of b) ___ hypotension
7. When cleaning if the site has significant drainage repeat the cleaning with a ___ alcohol swabstick
10. The ___ is responsible for identifying patients who are candidates for a whole blood lactat
15. In the ITA we use the sepsis protocol lactate ___ order
17. place the Statlock and excess catheter above the ___ fossa
18. Notify teh medical provider STAT to report lactate greater than ___ mmol/L using SBAR report

20. ___ is a clinical response arising from a nonspecific inflammatory insult
21. If the newly inserted PICC has blood oozing frin the exit site the dressing is to be changed within ___ hours
22. For a newly placed PICC with no blood at the exit site, dresing changes are to begin in one ___
23. Complete the order by signing 'per protocol' ___ cosign

Down

2. High lactate is an established indicator of tissue ___ in patients who are not yet ___ but are at risk for sever sepsis or septic shock
3. High lactate is an established indicator of tissue ___ in patients who are not yet ___ but are at risk for sever sepsis or septic shock
5. When cleaning the PICC site with alchol start at the insertions ite and clean in an outward motion to at least 4-5 ___ in diameter

8. measure the length of the PICC catheter extending from the exit sit to the ___ hub
9. During a dressing change you remove the old dressing in the ___ of the insertion
11. When perfomring dressing changes scrub hands with soap for minimum of ___ seconds
12. When not in use PICCs need to be flushed every ___ days
13. Chlorhexadine may take up to three ___ to dry
14. Apply the Biopatch with teh blue side up. Ensure the catheter lumens sit on ___ of the slit
16. The in-hospital mortality rate for patients with septicemia is ___ times higher than for other diagnoses
19. septic shock is severe sepsis with a) lactate > ___ or