
Name:	Date:
-------	-------

Sexually Transmitted Dieases



Across

- **2.** More common in woman then men, can cause discharge and premature delivery for women
- **5.** If you have intercourse, oral, anal and genital touching, you can get an
- **6.** interferes with your body's ability to fight off virus, bacteria and fungi that cause illness and lead to a chronic life threatening disease
- **9.** A bacterial infection of the genitals, anus or throat
- 10. How do you contract an STD?

- **11.** Often have no signs or symptoms, even with no symptoms you can still pass the infection to your sex partners
- **12.** Around 5% of women aged 14-24 will contract this, and that number may even be higher since this STI is difficult to identify. Left untreated can cause pelvic inflammatory disease.
- **14.** Usually easy to detect, blisters forming on or around the genitals tend to attract attention.
- **15.** What does STD Stand for? **Down**
- 1. What does STI stand for?

- **3.** A viral infection affecting the liver, mild illness lasting a short time
- **4.** Best protection from sexually transmitted diseases
- **7.** An infection caused by bacteria that can spread through out the body
- 8. Is a contagious liver disease that can range from mild illness to a serious life long condition. It can be transmitted sexually, but most often transmitted through contaminated needles
- **13.** Common in both men and women, but especially hazardous in certain cases to females