
Name:	Date:
i tairio.	Dato.

Shaken Baby Syndrome



Λ	_	na		
A	c	ra	22	

1. Fewer then	% of infants
who survive have normal	development

- 5. Thirty-five percent of victims of SBS are completely____ on their caregivers.
- 8. 50%-75% of ____ and adults surveryed did not know shaking an infant is dangerour
- 10. Violently chaking an infant can cause injury after as few as _____seconds.
- 11. Symptoms of SBS can start or peek within 4-6 hours.
- 12. Injury occurs when shaking an infant results in force that _____ the bridging veins between he dura mater and the brain substance.

14	The mortality	rate with	SRS	ic
		Tute with	363	13
appr	roximatelv	%		

- 15. It is _____ ok to shake an infant.
- 16. ____ is the soft spot on an infants head.
- 18. There are 1200 to 1600 reported cases of SBS in the _____ each year.
- 19. A long-term affect of SBS is
- 20. The degree of damage depends on the amount and _____ of shaking that child endured.

<u>Down</u>

2. Stepfathers and boyfriends cause 60% of SBS cases each year.

3. SBS ia a form of	3.	SBS	iα	a	form	of	
---------------------	----	-----	----	---	------	----	--

	
4	is
caused from violently shaking a	n infant
/ CDC	41

- 6. SBS causes more _____ then a fall from a 3 story building.
 7. Mothers are _____ in
- - 9. ____ is a long-term affect of SBS.
 - 13. SBS is most common in children under _____ years old.
 - 17. Children up to age ____ can be injuried by violent shaking.