


# Sleep, Parkinson's, Seizures



## Across

3. agents used for sedation, insomnia, anxiety, seizures  
 6. BUN and creatinine evaluate which organs  
 9. an effect of a chemical substance, opposite to the effect normally expected  
 14. substance to avoid prior to HS  
 15. agent produces sleep  
 16. succinimides, hydantoin, benzodiazepines and barbiturates-used to treat  
 19. antidote for benzodiazepine overdoses

21. appropriate method to discontinue a benzodiazepine  
 22. available in disintegrating tablets

## Down

1. Parkinson's patients have excess  
 2. serious adverse effect associated with dopamine agonists, pramipex and ropinirole  
 4. significant side effect of phenytoin  
 5. frequently used to treat status epilepticus  
 7. salivation, pill rolling, mask like facial expression are Parkinson's

8. foods high in this should be avoided if taking a MAOI  
 10. enhances the metabolism of benzodiazepines  
 11. agent that produces relaxation  
 12. vitamin b6, decreases the therapeutic effects of  
 13. benzodiazepines suggested duration of use  
 17. amantadine, levodopa, bromocriptine, used to treat  
 18. time frame for full therapeutic effect of levodopa (Larodopa)\_ months  
 20. AST, ALT, GGT, bilirubin, and platelet levels evaluate which organ