

Name: _____ Date: _____

sexually transmitted diseases

- | | |
|--|--------------------------------|
| 1. Avoid raw meat and cat litter as preventative management | A. HIV |
| 2. Infants will receive the HBIG at birth when the mother has this infection | B. Pelvic inflammatory disease |
| 3. This infection can lead to a 50 percent chance of deafness and blindness if exposed in the first trimester | C. 36 weeks |
| 4. no treatment for this asymptomatic and mononucleosis like manifestation | D. Trichomoniasis |
| 5. A bacterial infection that can be passed to the fetus during labor and delivery | E. GBS |
| 6. Administer retrovir from 14 weeks on and deliver by c/section gives a neonate the best chance to manage this STI | F. Rubella |
| 7. This STI is the most commonly reported STI among American women | G. Hepatitis B |
| 8. This medication is the most common medication used to treat chlamydia | H. Syphilis |
| 9. A vaginal and rectal culture to assess for GBS is typically done at this gestation | I. Gonorrhea |
| 10. Ceftriaxone IM and azithromycin together are usually given to treat this STI | J. Chlamydia |
| 11. Penicillin G IM in a single dose is usually given to treat this STI | K. Candidiasis |
| 12. Genital warts called condyloma acuminata that can lead to cervical cancer is associated with this Sexually transmitted virus | L. HPV human papilloma virus |
| 13. The STI caused by the protozoan parasite causes this STI | M. Toxoplasmosis |
| 14. Fluconazole is a medication that can be used in this overgrowth of yeast called | N. CMV |
| 15. This bacteria most commonly caused by haemophilus vaginalis or Gardnerella can lead to this | O. Azithromycin |