


Name: _____

TREATMENTS FOR SAH



ACROSS

- 2. WHAT DOES ICP STAND FOR
- 9. WHAT ANTICONVULSANT IS USUALLY GIVEN FOR SEIZURES
- 10. HOW MANY CRANIAL NERVES ARE THERE
- 11. WHAT SURGERY CAN BE PERFORMED TO SECURE AN ANEURYSM
- 13. HOW OFTEN DO YOU ADMINISTER NIMODIPINE

WORD BANK

- ABCDE APPROACH
- SAH
- FOUR HOURLY
- DECREASED CONSCIOUS LEVEL
- NIMODIPINE
- THREE LITRES

- 14. SMOKING IS A RISK FACTOR OF WHAT?
- 15. WHAT IS THE FIRST ASSESSMENT THAT SHOULD BE APPROACHED ON SOMEONE WITH REDUCED GCS
- 16. WHAT IS AN EVD USED FOR
- 17. WHAT IS THE RECOMMENDED FLUID INTAKE FOR SAH PATIENT

DOWN

- 1. WHAT MEDICATION DO WE ADMINISTER FOR 21 DAYS

- INTRACRANIAL PRESSURE
- GLASGOW COMA SCALE
- TWELVE
- SEVEN DAYS
- HYDROCEPHALUS
- LEVETIRACETAM

- 3. TYPICAL SYMPTOM OF SAH
- 4. WHAT MEDICATION IS A CALCIUM CHANNEL BLOCKER
- 5. SAH PATIENTS MAY HAVE...?
- 6. WHAT DOES SAH STAND FOR
- 7. GCS STANDS FOR
- 8. WHEN IS THE PEAK INCIDENCE FOR VASOSPASM
- 12. NAME THE TERM TO DESCRIBE WHEN BLOOD VESSELS CONSTRICT

- THUNDERCLAP HEADACHE
- VASOSPASM
- NIMODIPINE
- COILING
- SUBARACHNOID HEMORRHAGE